



Page 1:2

RIODINE S POVIDONE-IODINE SURGICAL SCRUB 100mg/mL

Composition

Povidone-Iodine 10%w/v (equivalent to 1%w/v available Iodine)

Actions

Povidone-lodine is an iodophore in which lodine is bound to the carrier molecule, Povidone. The complex, Povidone-lodine, slowly liberates inorganic lodine when in contact with the skin and mucous membranes. Only a small amount of lodine is released at any one time, giving Povidone-lodine a lower irritant potential, and longer duration of microbiocidal action than conventional lodine solutions.

Povidone-lodine is considered to have the same antimicrobial spectrum of lodine. It has broad-spectrum activity; including bacteria, fungi, viruses, protozoa and spores.

Indications

For preoperative and post operative scrubbing of skin by hospital operating personnel: for preoperative use on patients as an antiseptic cleanser or for general use as an antiseptic skin cleanser in the surgery.

Directions for Use

- 1. Use undiluted.
- 2. Wet hands and arms with warm water.
- 3. Use approximately 5mLon hands, wash for 2 minutes.
- 4. Use nailbrush for nails only.
- 5. Repeat wash before surgery.
- 6. Rinse well then dry carefully.

Contraindications

Known hypersensitivity to povidone-iodine or iodines.

Precautions

For external use only. Do not use near mucous membranes or open wounds. Prolonged exposure to wet solution may cause irritation or rarely, severe skin reaction. If skin irritation or rash occurs, discontinue use immediately.

Use in Pregnancy

lodine applied to intact skin is minimally absorbed (unless applied to a large enough surface area) however the benefits must be weighed against the risks. Possible effects of iodine absorption may include alteration of thyroid function of both mother and foetus.

Use in Lactation

See Use in Pregnancy.

Use in Children

Premature infants. Not recommended for use as an antiseptic scrub on premature newborn infants.

Full term infants. The use of Povidone-lodine containing preparations on full term infants may present a small risk of transient hypothyroidism especially if applied to large areas of the body and/or for extended periods of time (greater than five days). Repeated or large applications of Riodine S Povidone-lodine Surgical Scrub in full term infants should be used in conjunction with appropriate monitoring of thyroid function.

Adverse Effects

Very few adverse reactions have been reported from the use of povidone-iodine. Of these local irritation, itching and burning were most common. Should skin irritation occur, discontinue use.

First Aid

For advice, contact a doctor or **Poisons Information Centre (Phone Australia 131126; New Zealand 0800 764 766)**.

Storage Poison Schedule

Store below 25°C. Not Scheduled.

Presentation

Code	Product	Size
5000023708 (RIO00804F)	RIODINE S POVIDONE-IODINE SURGICAL SCRUB 100mg/mL (AUST R 21200)	500mL



USER GUIDE

Page 2:2

RIODINE S POVIDONE-IODINE SURGICAL SCRUB 100mg/mL

Contact Point

Perrigo Australia in Australia & Orion Laboratories (NZ) Ltd in New Zealand Orion Laboratories Pty Ltd, trading as Perrigo Australia ABN 56 009 293 136 25-29 Delawney Street, Balcatta, Western Australia 6021 AUSTRALIA

FREE PHONE: 1800 805 546 FREE FAX: 1800 004 110

EMAIL: customerservice@perrigo.com.au; WEBSITE: www.perrigo.com.au;

ORION® is a registered trademark of Orion Laboratories Pty Ltd

END OF USER GUIDE